

## ABUNDANCE AND PRODUCTIVITY ESTIMATES – 2023 UPDATE ATLANTIC COAST PIPING PLOVER POPULATION

Population monitoring on the breeding grounds has been an integral part of the recovery program for Atlantic Coast piping plovers since 1986, providing information to inform protection of breeding piping plovers and their habitat. Annual coastwide censuses are one component of monitoring that track local and regional progress toward recovery. This update describes the delisting criteria established in the recovery plan and discusses the role of abundance and distribution of breeding pairs in Atlantic Coast piping plover conservation. It summarizes the most current information about abundance and productivity with attention to changes since Endangered Species Act (ESA) listing and since the 2021 estimates that are reported in USFWS (2022).

### Recovery criteria and strategy

The objective of the 1996 revised Atlantic Coast Recovery Plan is to assure the long-term viability of the Atlantic Coast piping plover population in the wild, thereby allowing removal of this population from the Federal List of Endangered and Threatened Wildlife and Plants (50 CFR 17.11 and 17.12). The Atlantic Coast piping plover population may be considered for delisting when the following recovery criteria, established in the recovery plan, have been met:

1. Increase and maintain for 5 years a total of 2,000 breeding pairs, distributed among four recovery units.

| <u>Recovery Unit</u>                   | <u>Minimum Subpopulation</u> |
|----------------------------------------|------------------------------|
| Atlantic (Eastern) Canada <sup>1</sup> | 400 pairs                    |
| New England                            | 625 pairs                    |
| New York-New Jersey                    | 575 pairs                    |
| Southern (DE-MD-VA-NC)                 | 400 pairs                    |

2. Verify the adequacy of a 2,000-pair population of piping plovers to maintain heterozygosity and allelic diversity over the long term.
3. Achieve a 5-year average productivity of 1.5 fledged chicks per pair in each of the four recovery units described in criterion 1, based on data from sites that collectively support at least 90 percent of the recovery unit's population<sup>2</sup>.

---

<sup>1</sup> Canadian Wildlife Service documents and literature published since 2002 refer to piping plovers breeding in Nova Scotia, New Brunswick, Prince Edward Island, Quebec, and Newfoundland as the piping plover *melodus* subspecies or the "eastern Canada population." This subpopulation coincides exactly with the geographic area termed "Atlantic Canada Recovery Unit" in the Service's 1996 Recovery Plan. To reduce confusion, we refer henceforth in this status update to the Eastern Canada recovery unit.

<sup>2</sup> With regard to delisting criterion #3, the recovery plan further states that "The PVA [the population viability analysis, conducted to support development of the delisting criteria] shows that a population of only 2,000 pairs would remain highly vulnerable to extinction unless average productivity is sustained above 1.5 chicks per pair. However, since the PVA is based on several assumptions that *may* underestimate survival rates for some or all

4. Institute long-term agreements to assure protection and management sufficient to maintain the population targets and average productivity in each recovery unit.
5. Ensure long-term maintenance of wintering habitat sufficient in quantity, quality, and distribution to maintain survival rates needed for a 2,000-pair population.

The recovery strategy, as articulated in the plan, recognizes that attainment of abundance targets for each recovery unit increases the probability of survival and recovery of the entire population: “Dispersal of the population across its breeding range serves as a hedge against catastrophes, such as hurricanes, oil spills, or disease, which might depress regional survival and/or productivity. Maintaining robust, well-distributed subpopulations should reduce variance in survival and productivity of the Atlantic Coast population as a whole, facilitate interchange of genetic material between subpopulations, and promote recolonization of any sites that experience declines or local extirpations due to low productivity and/or temporary habitat succession (USFWS 1996).”

### **Role of Breeding Abundance and Distribution in Recovery**

As discussed below, the subpopulation abundance and distribution targets in recovery criterion #1 ensure **representation, redundancy, and resiliency** for Atlantic Coast piping plovers in their breeding range, consistent with current Service recovery planning guidance<sup>3</sup> (see also Schaffer and Stein 2000).

**Representation** supports the adaptability and evolutionary capacity of a species to accommodate long-term environmental changes (e.g., climate, habitat conditions or structure across large areas, emerging pathogens, novel competitors and/or predators, invasive species). The breadth of genetic, ecological, demographic, and behavioral diversity across a range of ecologically diverse locations or niches on the landscape are the best available and most useful expressions of representation (USFWS 2016b). A comprehensive molecular-genetic investigation of piping plovers by Miller et al. (2010) found strong genetic structure, supported by significant

---

recovery units and/or the percentage of one-year old adults that breed, this productivity figure may be revised downward if (1) it is demonstrated that survival rates are higher in some regions, and (2) a scientifically credible, stochastic model that incorporates the best available estimates of survival and other demographic variables shows that lower productivity rates will assure a 95 percent probability of survival for 100 years (see task 3.5). Adjustments to this criterion may be applied to the population as a whole or to one or more of the four recovery units, as supported by observed productivity and population trend data (USFWS 1996).” Citing findings of latitudinal variation in productivity needed to maintain a stationary population (Calvert et al. 2006, Hecht and Melvin 2009), the 2009 and 2020 Piping Plover 5-Year Reviews (USFWS 2009, 2020) recommends demographic modeling that explores effects of variation in productivity, survival rates, and carrying capacity of habitat on population viability within individual recovery units and the Atlantic Coast population as a whole to support revision of criterion #3.

<sup>3</sup> USFWS (2016a) states: “Recovery criteria: The objective, measurable thresholds for the parameters that contribute to the resiliency, redundancy, representation, including the level of amelioration of the factors negatively affecting the 3Rs (i.e., threats) needed to achieve the recovery vision for any species (delisting).”

correlations between genetic and geographic distances in both mitochondrial and microsatellite data sets for birds breeding along the Atlantic Coast from Newfoundland to North Carolina. Atlantic birds showed evidence of isolation-by-distance patterns, indicating that dispersal, when it occurs, is generally associated with movement to relatively proximal breeding territories. Maintaining geographically distributed subpopulations across the four recovery units serves to conserve representation of genetic diversity and adaptability to variable environmental selective pressures.

Analysis of a long-term data set found that both body mass of Atlantic Coast (and Northern Great Plains) piping plovers varied with latitude and environmental temperature in both their breeding and wintering ranges, while wing length correlated only with breeding range latitude and temperature. Differences in wing length are most likely driven by breeding season conditions or tradeoffs related to migration, a pattern that may have microevolutionary consequences (Gibson et al. 2019).

Further evidence of adaptive variability across recovery unit subpopulations is found in latitudinal differences in Atlantic Coast piping plover breeding habitat requirements. Zeigler et al (2021) found significant differences in the elevations, distances to ocean, and distances to low-energy shorelines of nesting habitats in the three U.S. Atlantic recovery units that are consistent with literature from more localized studies. Although piping plovers breeding in the northern part of their Atlantic Coast range mostly avoid sections of beach with high steep foredunes (Strauss 1990, Fraser et al. 2005), they are capable of thriving on beaches where chick access is limited to ocean foraging habitats<sup>4</sup> (Jones 1997, Boyne et al. 2014). In New York and New Jersey, however, the species demonstrates strong preference for sites that also offer chick access to ephemeral pools and bayside tidal flats (Elias et al. 2000, Cohen et al. 2009, Stantial et al. 2021, Robinson et al. 2021) and for flatter, sparsely vegetated habitats with moist substrates (Robinson et al. 2022). In Delaware, Maryland, Virginia, and North Carolina, Southern recovery unit breeding sites are almost completely restricted to low-lying barrier island flats and spits that also feature moist foraging substrates away from the ocean intertidal zone (McConnaughey et al. 1990, Loegering and Fraser 1995, Boettcher et al. 2007, NPS 2008). In addition to these well-documented geographic differences in habitat preferences, latitudinal variability may also provide some adaptive capacity for changing climatic factors such as breeding season temperatures and storm patterns that may affect Atlantic Coast piping plovers directly or indirectly (e.g., via changes in prey composition or phenology).

Another line of evidence for latitudinal adaptation within Atlantic Coast piping plovers is manifested in a strong pattern of higher productivity rates needed to maintain stable populations with increasing latitude (Hecht and Melvin 2009) and concomitant differences in annual survival rates. Although the underlying causes and mechanisms are not yet well understood, this striking demographic variability among recovery units may also contribute to evolutionary capacity. In summary, maintaining geographically well-distributed populations across the four recovery units serves to conserve representation of genetic diversity and adaptations to variable environmental selective pressures evidenced by genetic structure, diverse habitat requirements, and differences in vital rates.

---

<sup>4</sup> Management of human disturbance and human-abetted predation must also be provided.

**Redundancy** safeguards the ability of representative units to withstand catastrophic events. The number and distribution of resilient populations within each representative unit contribute to redundancy, thereby assuring that the loss of an individual population does not lead to loss of representation (USFWS 2016b). The 1996 recovery plan articulates the role of the recovery units in buffering Atlantic Coast piping plovers against catastrophic events such as large storms and oil spills during the breeding season, and this need is likewise served by attaining and maintaining robust, well-distributed populations within each recovery unit. Consistent with more qualitative characterization of high fidelity and low dispersal rates for Atlantic Coast piping plovers (USFWS 1996, Hecht and Melvin 2009, USFWS 2022), average interannual movement of breeding adults banded on southern Long Island between 2013 and 2022 was 2.8 kilometers (km; median 141.5 meters) and 88 percent were within 2.5 km of an individual's prior nest (Wails et al. 2023). Mean dispersal from natal sites to the first known nest was 20 km, and 75 percent were within 15 km (Wails et al. 2023). Two (0.6 percent) of 348 piping plovers banded in Eastern Canada in 2013-2018 and resighted in one or more subsequent breeding seasons nested in the U.S. Atlantic range (C. Gratto-Trevor, pers. comm. 2023). Thus, the ability of piping plovers in each recovery unit to rebound from events that depress productivity or survival and to colonize newly formed or improved habitat (e.g., after storms or artificial habitat enhancement projects) depends on redundancy that is measured via progress towards recovery unit abundance targets. Maintenance of abundance targets for at least 5 years provides evidence that recovery will be sustainable.

**Resiliency** is the ability to sustain populations in the face of demographic variation and environmental stochasticity. Resiliency depends on a number of vital rates that ultimately affect population size and growth rate, as well as distribution (USFWS 2016b). In the case of Atlantic Coast piping plovers, resiliency (like redundancy) is provided via widely distributed populations meeting abundance targets for breeding pairs within each recovery unit. Hecht and Melvin (2009) found significant positive relationships between productivity and population growth in the subsequent year for each of the three U.S. recovery units, and abundance of piping plovers in each recovery unit population is highly dependent on within-recovery unit productivity<sup>5</sup>. As noted above, dispersal rates decline steeply with distance from previous breeding and natal sites. Thus, robust numbers of evenly distributed breeding pairs support dispersal and within-recovery unit recolonization of any sites that experience declines or local extirpations due to low productivity and/or temporary habitat succession (Gilpin 1987, Goodman 1987, and Thomas 1994).

Wide distribution of breeding pairs within representative units also provides a buffer against environmental stochasticity. For example, weather events such as storms that flood nests may affect the south-facing beaches within a recovery unit in a given year more than north- and east-facing sites (or vice-versa). When environmental factors adversely affect productivity across a region, more abundant populations are inherently less susceptible to reaching the very low numbers from which it is difficult to rebound, and which make them vulnerable to local or regional extirpations if multiple years of poor productivity occur in close succession. Similarly, robust numbers of breeding pairs in each recovery unit will provide Atlantic Coast piping plovers with a buffer against stressors (e.g., weather, habitat degradation, disturbance) in their migration

---

<sup>5</sup> Maintaining survival (throughout the annual cycle) is also important.

and wintering range that may affect survival rates (Saunders et al. 2014, Gibson et al. 2018, Ellis et al. 2021).

Representation, redundancy, and resiliency are interconnected. Populations must be resilient in order to contribute to redundancy or representation. Likewise, redundant populations within a representative genotype or ecological setting contribute to maintenance of adaptive and evolutionary capacity (USFWS 2016b). For Atlantic Coast piping plovers, this is provided via subpopulation targets for four representative recovery units, thereby increasing the likelihood of survival and recovery of the Atlantic Coast population as a whole. Dispersal of the population across its breeding range in four robust subpopulations serves to protect against environmental and demographic variation and catastrophic events, and to conserve adaptive capacity.

## **Abundance and trends**

Abundance of Atlantic Coast piping plovers is reported as numbers of breeding pairs, that is, adult pairs that exhibit sustained ( $\geq 2$  weeks) territorial or courtship behavior at a site or are observed with nests or unfledged chicks (USFWS 1996). Annual estimates of breeding pairs of Atlantic Coast piping plovers are based on multiple surveys of almost all breeding habitat, including many currently unoccupied sites. Sites that cannot be monitored repeatedly in May and June (primarily sites with few pairs or inconsistent occupancy) are surveyed at least once during a standard 9-day count period (Hecht and Melvin 2009).

The 2023 Atlantic Coast piping plover population estimate of 2,593 pairs is 13 percent higher than the 2021 estimate and more than triple the estimate of pairs at the time of the 1986 ESA listing (Table 1). Discounting apparent increases in New York, New Jersey, and North Carolina between 1986 and 1989, which likely were due in part to increased census effort (USFWS 1996), the population increased 171 percent between 1989 and 2023.

Overall population growth is tempered by pronounced geographic and temporal variability (Figure 1). The largest population increase between 1989<sup>6</sup> and 2023 occurred in New England (648 percent). Abundance in the New York-New Jersey recovery unit experienced a net increase of 98 percent between 1989 and 2023. However, this population declined sharply from a peak of 586 pairs in 2007 to 378 pairs in 2014, before rebounding to 631 pairs in 2023. Net growth in the Southern recovery unit population was 17 percent between 1989 and 2023. Most of the Southern recovery unit breeding population increase occurred in 2003 to 2005 and 2011 to 2012, and the population decreased 40 percent between 2016 and 2023. In Eastern Canada, where increases have often been quickly eroded in subsequent years, the population posted a net 19-percent decline between 1989 and 2023. See further discussion of recovery unit trends below.

## **Productivity**

Atlantic Coast piping plover productivity is reported as number of chicks fledged per breeding pair. For purposes of measuring productivity, chicks are counted as fledged if they survive to 25 days of age or are seen flying, whichever occurs first. Productivity for each state and recovery

---

<sup>6</sup> As noted in the preceding paragraph, 1989 estimates provide the best baseline for assessing changes in abundance because they reflect increased survey effort in the three years following the ESA listing.

unit is calculated by dividing the number of fledged chicks by the number of pairs that were monitored and for which number of fledglings could be determined. This includes both successful pairs and pairs that fledged no chicks either because they failed to nest or because no eggs hatched or no chicks survived to fledging. Accurate assessment of productivity is facilitated by repeated visits to nesting beaches to monitor individual nests and broods during May, June, July, and, if necessary, August.

Hecht and Melvin (2009) found latitudinal trends in Atlantic Coast piping plover productivity and relationships between productivity and population growth. Average productivity from 1989 through 2006 decreased with decreasing latitude (Hecht and Melvin 2009). Within recovery units, productivity was variable from year to year and showed no sustained trends. There were significant, positive relationships between productivity and population growth in the subsequent year for each of the three U.S. recovery units, but not for Eastern Canada. Regression analysis indicated a latitudinal trend in predictions of annual productivity needed to support stationary populations within recovery units, increasing from 0.93 chicks fledged per pair in the Southern unit to 1.44 in Eastern Canada. Relatively small coefficients of determination ( $r^2 = 0.09$  to  $0.59$ ) for the relationships between annual productivity and population increases in the subsequent year indicate that other factors, most likely annual survival rates of both adults and fledged chicks, also had important influences on population growth rates. In some parts of the range, habitat availability may also be constraining recruitment into the breeding population.

Annual productivity estimates for the 1992-2023 period are summarized by recovery unit and state in Table 2. Average annual productivity for the U.S. Atlantic Coast during 1989-2023 was 1.24 fledged chicks per pair, and it continues to demonstrate a strong latitudinal trend, decreasing from north to south. The overall U.S. Atlantic Coast productivity estimate in 2022 was 1.17 fledged chicks per pair, 1.04 chicks per pair in 2023. Piping plovers in the Southern Recovery Unit experienced record-low productivity in both 2022 and 2023.

### **Status by Recovery Unit**

The demographic status of each recovery unit and implications for the survival and recovery of the coastwide population are summarized below.

**Eastern Canada recovery unit<sup>7</sup>** - Despite much higher long-term average productivity than the other recovery units, the Eastern Canada subpopulation decreased from 233 pairs in 1989 to 158 pairs in 2020 before rebounding to 189 pairs in 2023. Dependence of Eastern Canada breeding abundance on high prior-year productivity is illustrated by a decline (albeit by a single pair) in 2022 following productivity of 1.55 fledged chicks per pair in 2021. Breeding pairs increased 6

---

<sup>7</sup> The piping plover listing under the United States ESA designates the species as threatened across its range, except in the watershed of the Great Lakes where it is listed as endangered (USFWS 1985). The Committee on the Status of Endangered Wildlife in Canada (COSEWIC 2001) currently recognizes piping plovers breeding in Eastern Canada as *Charadrius melodus melodus* and designates the subspecies as “Endangered” (Department of Justice Canada 2002). This supersedes 1978 and 1985 designations assigned to the entire Canadian population of piping plovers (COSEWIC 2001). The Canadian piping plover recovery strategy recognizes the importance of conserving migration and wintering habitat (Environment Canada 2012). Canadian piping plover breeding sites identified as critical habitat receive legal protections under the Species at Risk Act (Environment Canada 2012).

percent in 2023 following productivity of 1.74 chicks per pair in 2022, very similar to Eastern Canada productivity of 1.76 chicks per pair in 2023.

In-depth evaluation of Eastern Canada piping plover population and productivity trends and environmental factors by the Wildlife Research Division of the Wildlife and Landscape Science Directorate, Environment Canada, concluded that the limiting factors now impeding recovery are primarily occurring outside Canada, during migration or on the wintering grounds (Gratto-Trevor et al. 2013). Efforts to identify these factors are in progress, but the difficulties inherent to discerning links between environmental factors in the nonbreeding range and vital demographic rates mean that rapid study results are unlikely. Furthermore, the availability of measures to ameliorate causal factors that may be identified is unknown. Meanwhile, Canadian Wildlife Service and other conservation partners continue ongoing intensive efforts to protect breeding habitat and activity to maximize productivity and reverse or slow the population decline. Low abundance, a 19 percent net population decline over the last 34 years (-29 percent since 2007), and lack of identified causal factors that can be remedied make the prospects for recovery of the Eastern Canada recovery unit highly uncertain.

**New England recovery unit** - The largest and most sustained population increase has occurred in New England, where the recovery unit population has exceeded (or been within 3 pairs of) its 625-pair abundance goal since 1998. The population posted a 7-percent increase in 2022, then grew 14 percent in 2023 to an estimated 1,541 pairs. Productivity in seven of the last ten years exceeded the estimated 1.21 chicks per pair needed to maintain a stationary population in New England (Hecht and Melvin 2009). Indeed, productivity of 1.56 fledged chicks per pair in 2019 matches or exceeds all productivity estimates attained since 1999. Productivity of 1.14 fledged chicks per pair in 2023 was below average but is not considered cause for concern in the context of current abundance of breeding pairs in New England.

Continuing growth of the New England piping plover population is generally attributable to the species' ability to breed productively on a wider range of microhabitats than in the other U.S. recovery units (Zeigler et al. 2021), along with intensive management of human disturbance and predation. Since 2016, the Massachusetts portion of the population has continued to increase concurrent with implementation of specific activities that may cause take of piping plovers under the auspices of the Massachusetts Habitat Conservation Plan (Massachusetts Division of Fisheries and Wildlife 2016). Additional carefully managed flexibility may be feasible for the relatively robust population in the New England recovery unit, especially if frameworks automatically adjust the amount of allowable flexibility in response to future changes in the abundance of breeding piping plovers<sup>8</sup>.

**New York-New Jersey recovery unit** – In 2023, abundance in the New York-New Jersey recovery unit reached a post-listing peak of 631 pairs, 10 percent above the 2021 breeding population and the recovery unit goal of 575 pairs. A prior post-listing peak of 586 pairs was attained in 2007, but the population began to decline the following year. Abundance of breeding

---

<sup>8</sup> For example, the Massachusetts Habitat Conservation Plan adjusts the amount of allowable flexible management in response to the size of the 3-year average breeding population in the State. This increases flexibility as the population increases, and it also provides a brake on authorized take in the event of a future decline in the breeding population (regardless of the cause).

pairs decreased 35 percent to 378 pairs in 2014 following 7 years of low productivity (including 4 years when it was less than 1.0 chick per pair). Improved productivity in 2014 and 2015 fueled a partial rebound to 496 pairs in 2016, but the population estimate increased by only one pair in 2017 and declined slightly in 2018 (despite high productivity in both 2016 and 2017) before increasing again in 2019. The apparent population dip in 2020 may be partially an artifact of decreased monitoring intensity in New York during the COVID-19 pandemic (McMaugh pers. comm. 2021). The recent population growth rate in New York-New Jersey has roughly tracked productivity in the preceding year, increasing 7 percent in 2022 following 2021 productivity of 1.19 chicks per pair, then increasing 2 percent in 2023 following productivity of 1.06 chicks per pair in 2022. Productivity in 2023 dipped to 1.03 chicks per pair.

Changes in the Long Island population account for most of the increases and decreases in the recovery unit population. The New Jersey piping plover population has fluctuated at low numbers (1989–2021 range = 92 to 144 pairs). In 2023, 74 percent of the New Jersey nesting pairs was concentrated on the Sandy Hook Unit of the Gateway National Recreation Area and the Holgate Unit of Edwin B. Forsythe National Wildlife Refuge (Heiser and Davis 2023), which comprise less than 8 percent of the State’s sandy ocean beaches in (Rice 2017). Low productivity of New Jersey piping plovers in 2021–2023 exacerbates concerns about the declining status of the adjacent Southern recovery unit.

Periodic declines and lack of population growth following some years with productivity well-above the rate estimated necessary to maintain a stationary population in the New York-New Jersey recovery unit (Hecht and Melvin 2009, Weithman et al. 2019) and response to habitat-creating events (Cohen et al. 2009, Robinson et al. 2019, 2020) suggest that habitat availability may be a strong factor affecting New York-New Jersey piping plover breeding abundance. Active habitat management to conserve the carrying capacity of breeding habitat in the New York-New Jersey recovery unit may be necessary for its piping plover population to sustain its abundance goal and thereby assure its long-term resiliency.

**Southern recovery unit** – The Southern recovery unit piping plover population decreased 14 percent between 2021 and 2023. Abundance of breeding pairs was 232 pairs, 40 percent lower than the post-listing high of 386 breeding pairs in 2016. Productivity rates in 2022 and 2023 (0.43 and 0.37 fledged chicks per pair, respectively) were the lowest on record, indicating that a further decline is very likely in 2024.

The Southern subpopulation responded positively to habitat creation events such as the 1992–1993 Nor’easters, Hurricane Isabel in 2003, Hurricane Ophelia in 2005, and Hurricane Irene in 2011 (Boettcher et al. 2007, Schupp et al. 2013, USFWS 2014, Robinson et al. 2019). Past years of low productivity (especially successive years of low productivity such as occurred in 2007–2008) have been followed by declines in breeding abundance, but the decline that began in 2016 is the steepest and most sustained that has been observed during the last 35 years (Figure 1). Ongoing collaborative efforts by Southern recovery unit biologists to identify and address the causes of such a large regional decline were initiated in early 2021, but no single explanatory factor has emerged to date. Increasing Southern recovery unit productivity and reversing the trend in breeding abundance are urgent priorities.

## Summary

Although population growth, from approximately 957 pairs in 1989 to an estimated 2,593 pairs in 2023, has reduced the Atlantic Coast piping plover's vulnerability to extinction since listing under the ESA, the distribution of population growth remains very uneven. Declines of 29 percent in the Eastern Canada breeding population since 2007 and 40 percent in the Southern recovery unit since 2016 typify long-standing concerns about the uneven distribution of Atlantic Coast piping plovers (Hecht and Melvin 2009, USFWS 2009, USFWS 2020). Future trends in breeding abundance will help inform assessments of whether current habitat and ongoing management are sufficient to sustain the New York-New Jersey subpopulation goal, attained in 2021. The New England recovery unit constitutes a stronghold, but there is no evidence of demographically meaningful dispersal to either Eastern Canada or New York-New Jersey, and any future inter-recovery unit "rescue" will be very slow. The survival and recovery of Atlantic Coast piping plovers remain dependent on rangewide conservation of remaining habitats and habitat-formation processes, as well as annual implementation of labor-intensive management to minimize the effects of pervasive and persistent threats from predation and disturbance by humans and pets (USFWS 2009, USFWS 2020).

## References Cited

- Boettcher, R., T. Penn, R.R. Cross, K.T. Terwilliger, and R.A. Beck. 2007. An overview of the status and distribution of piping plovers in Virginia. *Waterbirds* 30 (special publication 1):138-151.
- Boyne, A.W., D.L. Amirault\_Langlais, and A.J. McCue. 2014. Characteristics of piping plover nesting habitat in the Canadian Maritime Provinces. *Northeastern Naturalist*, 21(2):164-173.
- Calvert, A.M., D.L. Amirault, F. Shaffer, R. Elliot, A. Hanson, J. McKnight, and P.D. Taylor. 2006. Population assessment of an endangered shorebird: the Piping Plover (*Charadrius melodus*) in Eastern Canada. *Avian Conservation and Ecology* 1(3):4. Accessed on April 30, 2008, at <http://www.ace-eco.org/vol1/iss3/art4/>.
- Cohen, J.B., J.D. Fraser, and Daniel H. Catlin. 2006. Survival and site fidelity of piping plovers on Long Island, New York. *J. Field Ornithology*. 77(4):409-417.
- Cohen, J.B., L.M. Houghton, and J.D. Fraser. 2009. Nesting density and reproductive success of piping plovers in response to storm- and human- created habitat changes. *Wildlife Monographs* 173.
- COSEWIC. 2001. Canadian Species at Risk, May 2001. Committee on the Status of Endangered Wildlife Species in Canada. Ottawa, Ontario, Canada.
- Cross, R.R. (unpubl. data in USFWS 1996).
- Department of Justice Canada. 2002. Annual Statutes of Canada 2002, Chapter 29. Species at Risk Act, Schedule 1, Part 2.

- Elias, S.P., J.D. Fraser, and P.A. Buckley. 2000. Piping plover brood foraging ecology on New York barrier islands. *Journal of Wildlife Management*. 64(2): 346-354.
- Ellis, K.S., M.J. Anteau, F.J. Cuthbert, C.L. Gratto-Trevor, J.G. Jorgensen, D.J. Newstead, L.A. Powell, M.M. Ring, M.H. Sherfy, R.J. Swift, D.L. Toy and D.N. Koons. 2021. Impacts of extreme environmental disturbances on piping plover survival are partially moderated by migratory connectivity. *Biological Conservation* 264.
- Environment Canada. 2012. Recovery strategy for the piping plover (*Charadrius melodus melodus*) in Canada. Species at Risk Act Recovery Strategy Series. Environment Canada, Ottawa.
- Fraser, J.D., S.E. Keane, and P.A. Buckley. 2005. Pre-nesting use of intertidal habitats by piping plovers on South Monomoy Island, Massachusetts. *Journal of Wildlife Management* 69:1731-1736.
- Gibson, D., M.K. Chaplin, K.L. Hunt, M.J. Friedrich, C.E. Weithman, L.M. Addison, V. Cavalieri, S. Coleman, F.J. Cuthbert, J.D. Fraser, W. Golder, D. Hoffman, S.M. Karpanty, A. Van Zoeren, and D.H. Catlin. 2018. Impacts of anthropogenic disturbance on body condition, survival, and site fidelity of nonbreeding Piping Plovers. *The Condor* 120(3):566-580.
- Gibson, D., A. D. Hornsby, M. B. Brown, J. B. Cohen, L. R. Dinan, J. D. Fraser, M. J. Friedrich, C. L. Gratto-Trevor, K. L. Hunt, M. Jeffery, J. G. Jorgensen, P. W. C. Paton, S. G. Robinson, J. Rock, M. L. Stantial, C. E. Weithman and D. H. Catlin. 2019. Migratory shorebird adheres to Bergmann's Rule by responding to environmental conditions through the annual lifecycle. *Ecography* 42:1-12. doi: 10.1111/ecog.04325.
- Gilpin, M.E. 1987. Spatial structure and population vulnerability, pp 125-139 in M.E. Soule (ed) *Viable populations for Conservation*. Cambridge University Press, New York.
- Goodman, D. 1987. How do species persist? Lessons for Conservation Biology. *Conservation Biology* 1: 59-62.
- Gratto-Trevor, C.L., G.J. Robertson, and C.A. Bishop. 2013. Scientific review of the recovery program for Piping Plover (*melodus* subspecies) in Eastern Canada. Unpublished report, Science and Technology Branch, Environment Canada, PNWRC, 115 Perimeter Road, Saskatoon, Saskatchewan.
- Hecht, A., and S.M. Melvin. 2009. Population trends of Atlantic Coast piping plovers, 1986-2006. *Waterbirds* 32:64-72.
- Heiser, E. and C. Davis. 2023. Piping plover nesting results in New Jersey: 2023. New Jersey Department of Environmental Protection, Fish and Wildlife, Endangered and Nongame Species Program.

- Jones, K. 1997. Piping plover habitat selection, home range, and reproductive success at Cape Cod National Seashore, Massachusetts. National Park Service Technical Report NPS/NESO-RNR/NRTR/97-03.
- Loefering, J.P. 1992. Piping plover breeding biology, foraging ecology, and behavior on Assateague Island National Seashore, Maryland. M.S. Thesis. Virginia Polytechnic Institute and State University, Blacksburg, Virginia.
- Loefering, J.P. and J.D. Fraser. 1995. Factors affecting piping plover chick survival in different brood-rearing habitats. *Journal of Wildlife Management* 59(4): 646-655.
- MacIvor, L.H., C.R. Griffin, and S.M. Melvin. 1987. Management, habitat selection, and population dynamics of piping plovers on outer Cape Cod Massachusetts; 1985-1987. Submitted to National Park Service, Cape Cod National Seashore, South Wellfleet, Massachusetts.
- Massachusetts Division of Fisheries and Wildlife. 2016. Massachusetts Division of Fisheries & Wildlife (DFW) habitat conservation plan for piping plover. Prepared by MADFW and ICF International. Westborough, Massachusetts and Fairfax, Virginia.
- Massachusetts Division of Fisheries and Wildlife. 2022. Summary of the 2021 Massachusetts piping plover census. Natural Heritage & Endangered Species Program, Massachusetts Division of Fisheries & Wildlife. February 2022.
- McConnaughey, J.L., J.D. Fraser, S.D. Coutu, and J.P. Loefering. 1990. Piping plover distribution and reproductive success on Cape Lookout National Seashore. Unpublished Report submitted to the National Park Service. Virginia Polytechnic Institute and State University, Blacksburg, Virginia.
- Miller, M.P., S.M. Haig, C.L. Gratto-Trevor, and T.D. Mullins. 2010. Subspecies status and population genetic structure in piping plover (*Charadrius melodus*). *Auk* 127:57-71.
- National Park Service [NPS]. 2008. Piping plover (*Charadrius melodus*) monitoring at Cape Lookout National Seashore, 2008 summary report. Cape Lookout National Seashore, Harkers Island, North Carolina.
- New Jersey Division of Fish and Wildlife. 2021. Piping plover nesting results in New Jersey: 2021. New Jersey Division of Fish and Wildlife, Endangered Species and Nongame Species Program.
- Rice, T.M. 2016. Inventory of habitat modifications to tidal inlets in the U.S. Atlantic Coast breeding range of the piping plover (*Charadrius melodus*) as of 2015: Maine to North Carolina. Report submitted to the U.S. Fish and Wildlife Service, Hadley, Massachusetts.
- Rice, T.M. 2017. Inventory of habitat modifications to sandy oceanfront beaches in the U.S. Atlantic Coast breeding range of the piping plover (*Charadrius melodus*) as of 2015: Maine to North Carolina. Report submitted to the U.S. Fish and Wildlife Service, Hadley, Massachusetts.

- Rioux, S., D.L. Amirault-Langlais, and F. Shaffer. 2011. Piping plover make decisions regarding dispersal based on personal and public information in a variable coastal ecosystem. *Journal of Field Ornithology* 82:32–43.
- Robinson, S.G., J. Fraser, D. Catlin, S.M. Karpanty, J. Altman, R. Boettcher, K. Holcomb, C. Huber, K. Hunt, and A. Wilke. 2019. Irruptions: evidence for breeding season habitat limitation in piping plover (*Charadrius melodus*). *Avian Conservation and Ecology* 14(1):19. <https://doi.org/10.5751/ACE-01373-140119>.
- Robinson, S.G., D. Gibson, T. V. Riecke, J. D. Fraser, H. A. Bellman, A. DeRose-Wilson, S.M. Karpanty, K. M. Walker, and D. H. Catlin. 2020. Piping plover population increase after Hurricane Sandy mediated by immigration and reproductive output. *The Condor*:122. DOI: 10.1093/condor/duaa041.
- Robinson, S., H. Bellman, K. Walker, D. Catlin, S. Karpanty, S. Ritter, and J. Fraser. 2021. Adult piping plover habitat selection varies by behavior. *Ecosphere* 12(12):e03870. 10.1002/ecs2.3870.
- Robinson, S. G., K. M. Walker, H. A. Bellman, D. Gibson, D. H. Catlin, S. M. Karpanty, S. J. Ritter, and J. D. Fraser. 2022. Piping plover chick ecology following landscape-level disturbance. *Journal of Wildlife Management* e22325. <https://doi.org/10.1002/jwmg.22325>.
- Saunders, S.P., T.W. Arnold, E.A. Roche, and F.J. Cuthbert. 2014. Age-specific recruitment of piping plovers *Charadrius melodus* in the Great Lakes region. *Journal of Avian Biology*, 45(5) 437–449.
- Schaffer, M.L. and B.A. Stein. 2000. Safeguarding our precious heritage. Pages 301–321 in B.A. Stein, L.S. Kutner, and J.S. Adams, eds. *Precious heritage: The status of biodiversity in the United States*. Oxford University Press.
- Schupp, C.A., N.T. Winn, T.L. Pearl, J.P. Kumer, T.J.B. Carruthers, and C.S. Zimmerman. 2013. Restoration of overwash processes creates piping plover (*Charadrius melodus*) habitat on a barrier island (Assateague Island, Maryland). *Estuarine, Coastal, and Shelf Science* 116:11-20.
- Stantial, M.L., J.B. Cohen, A.J. Darrah, S. Farrell, and B. Maslo. 2021. Habitat-specific behavior, growth rate, and survival of piping plover chicks in New Jersey, USA. *Ecosphere* 12(10):e03782. 10.1002/ecs2.3782.
- Strauss, E. 1990. Reproductive success, life history patterns, and behavioral variation in populations of piping plovers subjected to human disturbance (1982-1989). Ph.D. Dissertation. Tufts University, Medford, MA.
- Thomas, C.D. 1994. Extinction, colonization, and metapopulations: environmental tracking by rare species. *Conservation Biology* 8:373-378.

- U.S. Fish and Wildlife Service [USFWS]. 1985. Determination of endangered and threatened status for the piping plover. Federal Register 50:50726-50734.
- USFWS. 1996. Piping plover (*Charadrius melodus*), Atlantic Coast population, revised recovery plan. Hadley, Massachusetts.
- USFWS. 2009. Piping plover (*Charadrius melodus*) 5-year review: Summary and evaluation. Northeast Regional Office, Hadley, Massachusetts.
- USFWS. 2014. Biological Opinion and Conference Opinion Fire Island Inlet to Moriches Inlet Fire Island Stabilization Project, Suffolk County, New York. Prepared for U.S. Army Corps of Engineers, New York District by U.S. Fish and Wildlife Service, Northeast Region. Dated October 15, 2014.
- USFWS. 2016a. Terms for recovery planning and implementation REV (Recovery Enhancement Vision) Training Class. September 2016.
- USFWS. 2016b. Clarification of the 3R's for the purpose of identifying recovery criteria. FWS, August 2016.
- USFWS. 2020. Piping plover (*Charadrius melodus*) 5-year review: Summary and evaluation. Michigan Field Office, East Lansing, Michigan and Northeast Regional Office, Hadley, Massachusetts.
- U.S. Fish and Wildlife Service. 2022. Abundance and productivity estimates – 2021 update: Atlantic Coast piping plover population. Hadley, Massachusetts.
- Wails, C. N., H. L. VanDerwater, H. A. Bellman, K. M. Black, A. DeRose-Wilson, S. S. Dorsey, K. W. Oliver, S. J. Ritter, S. G. Robinson, D. H. Catlin, S. M. Karpanty, and J. D. Fraser. 2023. Effectiveness monitoring and targeted studies in support of FIMI conservation measures. Annual report prepared by the Virginia Tech Shorebird Project. Virginia Tech, Blacksburg, Virginia.
- Weithman, C. E., S.G. Robinson, K.L. Hunt, J. Altman, H.A. Bellman, A.L. DeRose-Wilson, K.M. Walker, J.D. Fraser, S.M. Karpanty, and D.H. Catlin. 2019. Growth of two Atlantic Coast piping plover populations. The Condor 121. DOI: 10.1093/condor/duz037.
- Wilcox, L. 1959. A twenty year banding study of the piping plover. Auk 76:129-152.
- Zeigler, S.L., B.T. Gutierrez, A. Hecht, N.G. Plant, and E.J. Sturdivant. 2021. Piping plovers demonstrate regional differences in nesting habitat selection patterns along the U.S. Atlantic coast. Ecosphere 12(3):e03418.

***Electronic and personal communications***

Gratto-Trevor, C.L. 2023. Environment and Climate Change Canada. 01 December 2023 email to Anne Hecht, USFWS Northeast Region.

McMaugh, A. June 9, 2021 email from Andria McMaugh, New York State Department of Environmental Conservation to Anne Hecht regarding 2020 piping plover estimates.

***Recommended citation:*** U.S. Fish and Wildlife Service. 2024. Abundance and productivity estimates – 2023 update: Atlantic Coast piping plover population. Hadley, Massachusetts.

Table 1. Estimated abundance of Atlantic Coast piping plovers 1986-2023.

| State/RECOVERY<br>UNIT         | Pairs |      |      |      |      |      |      |      |      |      |      |      |      |      |      |      |      |      |      |      |      |
|--------------------------------|-------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|
|                                | 1986  | 1987 | 1988 | 1989 | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | 2006 |
| Maine                          | 15    | 12   | 20   | 16   | 17   | 18   | 24   | 32   | 35   | 40   | 60   | 47   | 60   | 56   | 50   | 55   | 66   | 61   | 55   | 49   | 40   |
| New Hampshire                  |       |      |      |      |      |      |      |      |      |      |      | 5    | 5    | 6    | 6    | 7    | 7    | 7    | 4    | 3    | 3    |
| Massachusetts                  | 139   | 126  | 134  | 137  | 140  | 160  | 213  | 289  | 352  | 441  | 454  | 483  | 495  | 501  | 496  | 495  | 538  | 511  | 488  | 467  | 482  |
| Rhode Island                   | 10    | 17   | 19   | 19   | 28   | 26   | 20   | 31   | 32   | 40   | 50   | 51   | 46   | 39   | 49   | 52   | 58   | 71   | 70   | 69   | 72   |
| Connecticut                    | 20    | 24   | 27   | 34   | 43   | 36   | 40   | 24   | 30   | 31   | 26   | 26   | 21   | 22   | 22   | 32   | 31   | 37   | 40   | 34   | 37   |
| NEW ENGLAND                    | 184   | 179  | 200  | 206  | 228  | 240  | 297  | 376  | 449  | 552  | 590  | 612  | 627  | 624  | 623  | 641  | 700  | 687  | 657  | 622  | 634  |
| New York                       | 106   | 135  | 172  | 191  | 197  | 191  | 187  | 193  | 209  | 249  | 256  | 256  | 245  | 243  | 289  | 309  | 369  | 386  | 384  | 374  | 422  |
| New Jersey                     | 102   | 93   | 105  | 128  | 126  | 126  | 134  | 127  | 124  | 132  | 127  | 115  | 93   | 107  | 112  | 122  | 138  | 144  | 135  | 111  | 116  |
| NY-NJ                          | 208   | 228  | 277  | 319  | 323  | 317  | 321  | 320  | 333  | 381  | 383  | 371  | 338  | 350  | 401  | 431  | 507  | 530  | 519  | 485  | 538  |
| Delaware                       | 8     | 7    | 3    | 3    | 6    | 5    | 2    | 2    | 4    | 5    | 6    | 4    | 6    | 4    | 3    | 6    | 6    | 6    | 7    | 8    | 9    |
| Maryland                       | 17    | 23   | 25   | 20   | 14   | 17   | 24   | 19   | 32   | 44   | 61   | 60   | 56   | 58   | 60   | 60   | 60   | 59   | 66   | 63   | 64   |
| Virginia                       | 100   | 100  | 103  | 121  | 125  | 131  | 97   | 106  | 96   | 118  | 87   | 88   | 95   | 89   | 96   | 119  | 120  | 114  | 152  | 192  | 202  |
| North Carolina                 | 30    | 30   | 40   | 55   | 55   | 40   | 49   | 53   | 54   | 50   | 35   | 52   | 46   | 31   | 24   | 23   | 23   | 24   | 20   | 37   | 46   |
| South Carolina                 | 3     |      | 0    |      | 1    | 1    |      | 1    |      |      | 0    |      |      |      |      | 0    |      |      |      |      |      |
| SOUTHERN                       | 158   | 160  | 171  | 199  | 201  | 194  | 172  | 181  | 186  | 217  | 189  | 204  | 203  | 182  | 183  | 208  | 209  | 203  | 245  | 300  | 321  |
| U.S. TOTAL                     | 550   | 567  | 648  | 724  | 752  | 751  | 790  | 877  | 968  | 1150 | 1162 | 1187 | 1168 | 1156 | 1207 | 1280 | 1416 | 1420 | 1421 | 1407 | 1493 |
| EASTERN<br>CANADA <sup>a</sup> | 240   | 223  | 238  | 233  | 230  | 252  | 223  | 223  | 194  | 200  | 202  | 199  | 211  | 236  | 230  | 250  | 274  | 256  | 237  | 217  | 256  |
| ATLANTIC<br>COAST TOTAL        | 790   | 790  | 886  | 957  | 982  | 1003 | 1013 | 1100 | 1162 | 1350 | 1364 | 1386 | 1379 | 1392 | 1437 | 1530 | 1690 | 1676 | 1658 | 1624 | 1749 |

<sup>a</sup> Includes 0-5 pairs on the French Islands of St. Pierre and Miquelon, reported by Canadian Wildlife Service.

Table 1 (continued). Estimated abundance of Atlantic Coast piping plovers 1986-2023.

| State/RECOVERY<br>UNIT         | Pairs |      |      |      |      |      |      |      |      |      |      |      |      |      |      |      |      |
|--------------------------------|-------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|
|                                | 2007  | 2008 | 2009 | 2010 | 2011 | 2012 | 2013 | 2014 | 2015 | 2016 | 2017 | 2018 | 2019 | 2020 | 2021 | 2022 | 2023 |
| Maine                          | 35    | 24   | 27   | 30   | 33   | 42   | 44   | 50   | 62   | 66   | 64   | 68   | 89   | 98   | 125  | 140  | 157  |
| New Hampshire                  | 3     | 3    | 5    | 4    | 4    | 6    | 7    | 6    | 8    | 7    | 7    | 9    | 11   | 12   | 13   | 14   | 16   |
| Massachusetts                  | 558   | 566  | 593  | 591  | 656  | 676  | 666  | 663  | 683  | 641  | 650  | 688  | 743  | 794  | 967  | 1033 | 1178 |
| Rhode Island                   | 73    | 77   | 84   | 85   | 86   | 90   | 92   | 91   | 99   | 97   | 87   | 87   | 80   | 85   | 99   | 99   | 111  |
| Connecticut                    | 36    | 41   | 44   | 43   | 52   | 51   | 45   | 51   | 62   | 63   | 66   | 64   | 57   | 58   | 60   | 66   | 79   |
| NEW ENGLAND                    | 705   | 711  | 753  | 753  | 831  | 865  | 854  | 861  | 914  | 874  | 874  | 916  | 980  | 1047 | 1264 | 1352 | 1541 |
| New York                       | 457   | 443  | 437  | 390  | 318  | 342  | 289  | 286  | 308  | 381  | 392  | 390  | 426  | 405  | 439  | 500  | 513  |
| New Jersey                     | 129   | 111  | 105  | 108  | 111  | 121  | 108  | 92   | 108  | 115  | 105  | 96   | 114  | 103  | 137  | 118  | 118  |
| NY-NJ                          | 586   | 554  | 542  | 498  | 429  | 463  | 397  | 378  | 416  | 496  | 497  | 486  | 540  | 508  | 576  | 618  | 631  |
| Delaware                       | 9     | 10   | 10   | 9    | 8    | 7    | 6    | 6    | 6    | 8    | 13   | 16   | 19   | 21   | 24   | 24   | 31   |
| Maryland                       | 64    | 49   | 45   | 44   | 36   | 41   | 45   | 38   | 36   | 34   | 34   | 23   | 24   | 24   | 22   | 19   | 8    |
| Virginia                       | 199   | 208  | 193  | 192  | 188  | 259  | 251  | 245  | 256  | 291  | 269  | 227  | 233  | 201  | 183  | 162  | 148  |
| North Carolina                 | 61    | 64   | 54   | 61   | 62   | 70   | 56   | 65   | 64   | 53   | 43   | 29   | 33   | 31   | 40   | 36   | 45   |
| South Carolina                 | 0     |      |      |      |      |      |      |      |      |      |      |      |      |      |      |      |      |
| SOUTHERN                       | 333   | 331  | 302  | 306  | 294  | 377  | 358  | 354  | 362  | 386  | 359  | 295  | 309  | 277  | 269  | 241  | 232  |
| U.S. TOTAL                     | 1624  | 1596 | 1597 | 1557 | 1554 | 1705 | 1609 | 1593 | 1692 | 1756 | 1730 | 1697 | 1829 | 1832 | 2109 | 2211 | 2404 |
| EASTERN<br>CANADA <sup>a</sup> | 266   | 253  | 252  | 225  | 209  | 179  | 184  | 186  | 179  | 176  | 173  | 181  | 190  | 158  | 180  | 179  | 189  |
| ATLANTIC<br>COAST TOTAL        | 1890  | 1849 | 1849 | 1782 | 1763 | 1884 | 1793 | 1779 | 1871 | 1932 | 1903 | 1878 | 2019 | 1990 | 2289 | 2390 | 2593 |

<sup>a</sup> Includes 0-5 pairs on the French Islands of St. Pierre and Miquelon, reported by Canadian Wildlife Service.

Table 2. Estimated productivity of Atlantic Coast piping plovers 1987-2023.

| State/RECOVERY<br>UNIT <sup>a</sup> | Chicks fledged per pair |      |      |      |      |      |      |      |      |      |      |      |      |      |      |      |      |      |      |      |
|-------------------------------------|-------------------------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|
|                                     | 1987                    | 1988 | 1989 | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | 2006 |
| Maine                               | 1.75                    | 0.75 | 2.38 | 1.53 | 2.50 | 2.00 | 2.38 | 2.00 | 2.38 | 1.63 | 1.98 | 1.47 | 1.63 | 1.60 | 1.98 | 1.39 | 1.28 | 1.45 | 0.55 | 1.35 |
| New Hampshire                       |                         |      |      |      |      |      |      |      |      |      | 0.60 | 2.40 | 2.67 | 2.33 | 2.14 | 0.14 | 1.00 | 1.00 | 0.00 | 0.67 |
| Massachusetts                       | 1.10                    | 1.29 | 1.59 | 1.38 | 1.72 | 2.03 | 1.92 | 1.81 | 1.62 | 1.35 | 1.33 | 1.50 | 1.60 | 1.09 | 1.49 | 1.14 | 1.26 | 1.38 | 1.14 | 1.33 |
| Rhode Island                        | 1.12                    | 1.58 | 1.47 | 0.88 | 0.77 | 1.55 | 1.80 | 2.00 | 1.68 | 1.56 | 1.34 | 1.13 | 1.79 | 1.20 | 1.50 | 1.95 | 1.03 | 1.50 | 1.43 | 1.03 |
| Connecticut                         | 1.29                    | 1.70 | 1.79 | 1.63 | 1.39 | 1.45 | 0.38 | 1.47 | 1.35 | 1.31 | 1.69 | 1.05 | 1.45 | 1.86 | 1.22 | 1.87 | 1.30 | 1.35 | 1.62 | 2.14 |
| NEW ENGLAND                         | 1.19                    | 1.32 | 1.68 | 1.38 | 1.62 | 1.91 | 1.85 | 1.81 | 1.67 | 1.40 | 1.39 | 1.46 | 1.62 | 1.18 | 1.53 | 1.26 | 1.24 | 1.40 | 1.15 | 1.34 |
| New York                            | 0.90                    | 1.24 | 1.02 | 0.80 | 1.09 | 0.98 | 1.24 | 1.34 | 0.97 | 1.14 | 1.36 | 1.09 | 1.35 | 1.11 | 1.27 | 1.62 | 1.15 | 1.46 | 1.44 | 1.55 |
| New Jersey                          | 0.85                    | 0.94 | 1.12 | 0.93 | 0.98 | 1.07 | 0.93 | 1.16 | 0.98 | 1.00 | 0.39 | 1.09 | 1.34 | 1.40 | 1.29 | 1.17 | 0.92 | 0.61 | 0.77 | 0.84 |
| NY-NJ                               | 0.86                    | 1.03 | 1.08 | 0.88 | 1.04 | 1.02 | 1.08 | 1.25 | 0.97 | 1.07 | 1.02 | 1.09 | 1.35 | 1.19 | 1.28 | 1.49 | 1.07 | 1.23 | 1.28 | 1.36 |
| Delaware                            |                         | 0.00 | 2.33 | 2.00 | 1.60 | 1.00 | 0.50 | 2.50 | 2.00 | 0.50 | 1.00 | 0.83 | 1.50 | 1.67 | 1.50 | 1.17 | 2.33 | 1.14 | 1.50 | 1.44 |
| Maryland                            | 1.17                    | 0.52 | 0.90 | 0.79 | 0.41 | 1.00 | 1.79 | 2.41 | 1.73 | 1.49 | 1.02 | 1.30 | 1.09 | 0.80 | 0.92 | 1.85 | 1.56 | 1.86 | 1.25 | 1.06 |
| Virginia                            |                         | 1.02 | 1.16 | 0.65 | 0.88 | 0.59 | 1.45 | 1.66 | 1.00 | 1.54 | 0.71 | 1.01 | 1.21 | 1.42 | 1.52 | 1.19 | 1.90 | 2.23 | 1.52 | 1.19 |
| North Carolina                      |                         |      | 0.59 | 0.43 | 0.07 | 0.41 | 0.74 | 0.36 | 0.45 | 0.86 | 0.23 | 0.61 | 0.48 | 0.54 | 0.50 | 0.17 | 0.46 | 0.65 | 0.92 | 0.87 |
| SOUTHERN                            | 1.17                    | 0.85 | 0.88 | 0.72 | 0.68 | 0.62 | 1.18 | 1.37 | 1.05 | 1.34 | 0.68 | 0.99 | 1.04 | 1.09 | 1.22 | 1.27 | 1.63 | 1.95 | 1.38 | 1.12 |
| U.S. average                        | 1.04                    | 1.11 | 1.28 | 1.06 | 1.22 | 1.35 | 1.47 | 1.56 | 1.35 | 1.30 | 1.16 | 1.27 | 1.45 | 1.17 | 1.40 | 1.34 | 1.24 | 1.43 | 1.24 | 1.30 |
| EASTERN<br>CANADA <sup>b</sup>      |                         | 1.65 | 1.58 | 1.62 | 1.07 | 1.55 | 0.69 | 1.25 | 1.69 | 1.72 | 2.10 | 1.84 | 1.74 | 1.47 | 1.77 | 1.18 | 1.62 | 1.93 | 1.82 | 1.82 |

<sup>a</sup> Annual recovery unit productivity is an average for all pairs breeding in the recovery unit in the pertinent year.

<sup>b</sup> Includes productivity on the French islands of St. Pierre and Miquelon, reported by Canadian Wildlife Service.

Table 2 (continued). Estimated productivity of Atlantic Coast piping plovers 1987-2023.

| State/RECOVERY<br>UNIT <sup>a</sup> | Chicks fledged per pair |      |      |      |      |      |      |      |      |      |      |      |      |      |      |      |      |                      |
|-------------------------------------|-------------------------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|----------------------|
|                                     | 2007                    | 2008 | 2009 | 2010 | 2011 | 2012 | 2013 | 2014 | 2015 | 2016 | 2017 | 2018 | 2019 | 2020 | 2021 | 2022 | 2023 | Average <sup>b</sup> |
| Maine                               | 1.06                    | 1.75 | 1.70 | 1.63 | 2.12 | 1.52 | 1.93 | 1.94 | 1.95 | 1.53 | 1.59 | 1.88 | 1.97 | 2.03 | 1.70 | 1.80 | 1.28 | 1.74                 |
| New Hampshire                       | 0.33                    | 2.00 | 0.40 | 1.50 | 2.00 | 0.67 | 1.71 | 0.33 | 1.50 | 2.14 | 0.71 | 1.89 | 1.82 | 1.33 | 1.00 | 2.00 | 2.38 | 1.36                 |
| Massachusetts                       | 1.25                    | 1.41 | 0.91 | 1.50 | 1.18 | 0.85 | 0.87 | 1.18 | 1.30 | 1.46 | 1.08 | 1.30 | 1.54 | 1.31 | 1.06 | 1.31 | 1.11 | 1.35                 |
| Rhode Island                        | 1.48                    | 1.68 | 1.46 | 1.76 | 1.49 | 1.06 | 0.98 | 1.63 | 1.58 | 1.49 | 0.66 | 0.87 | 1.20 | 1.41 | 1.23 | 0.95 | 0.86 | 1.35                 |
| Connecticut                         | 1.92                    | 2.49 | 1.68 | 1.91 | 1.37 | 1.18 | 1.82 | 2.27 | 1.81 | 1.38 | 1.52 | 1.17 | 1.72 | 1.00 | 1.33 | 1.47 | 1.35 | 1.53                 |
| NEW ENGLAND                         | 1.30                    | 1.51 | 1.04 | 1.56 | 1.27 | 0.93 | 1.00 | 1.33 | 1.41 | 1.47 | 1.10 | 1.30 | 1.56 | 1.37 | 1.15 | 1.35 | 1.14 | 1.39                 |
| New York                            | 1.15                    | 1.21 | 0.93 | 0.79 | 1.07 | 0.72 | 0.71 | 1.30 | 1.52 | 1.72 | 1.32 | 1.47 | 1.36 | 1.41 | 1.30 | 1.11 | 1.13 | 1.21                 |
| New Jersey                          | 0.67                    | 0.64 | 1.05 | 1.39 | 1.18 | 0.72 | 0.85 | 1.36 | 1.29 | 1.35 | 1.29 | 1.51 | 1.24 | 1.29 | 0.85 | 0.85 | 0.53 | 1.03                 |
| NY-NJ                               | 1.03                    | 1.10 | 0.96 | 0.92 | 1.09 | 0.72 | 0.74 | 1.32 | 1.46 | 1.62 | 1.32 | 1.48 | 1.33 | 1.38 | 1.19 | 1.06 | 1.03 | 1.16                 |
| Delaware                            | 1.33                    | 0.30 | 1.30 | 1.56 | 1.00 | 1.00 | 1.17 | 1.33 | 1.17 | 1.63 | 1.08 | 2.25 | 2.74 | 2.24 | 0.79 | 1.42 | 0.94 | 1.42                 |
| Maryland                            | 0.78                    | 0.41 | 1.42 | 1.09 | 1.25 | 1.02 | 0.76 | 1.55 | 1.31 | 1.47 | 0.82 | 1.30 | 1.00 | 0.21 | 0.68 | 0.53 | 0.50 | 1.12                 |
| Virginia                            | 1.16                    | 0.87 | 1.19 | 1.35 | 1.36 | 0.95 | 1.15 | 1.34 | 1.26 | 0.92 | 0.68 | 0.76 | 0.72 | 0.39 | 0.52 | 0.30 | 0.31 | 1.09                 |
| North Carolina                      | 0.26                    | 0.30 | 0.70 | 0.77 | 0.77 | 0.59 | 0.96 | 0.22 | 0.64 | 0.15 | 0.26 | 0.90 | 0.61 | 0.52 | 0.43 | 0.25 | 0.11 | 0.51                 |
| SOUTHERN                            | 0.92                    | 0.67 | 1.14 | 1.20 | 1.21 | 0.89 | 1.07 | 1.15 | 1.15 | 0.88 | 0.65 | 0.90 | 0.86 | 0.54 | 0.54 | 0.43 | 0.37 | 0.99                 |
| U.S. average                        | 1.13                    | 1.19 | 1.03 | 1.27 | 1.21 | 0.86 | 0.94 | 1.29 | 1.37 | 1.38 | 1.08 | 1.28 | 1.38 | 1.25 | 1.09 | 1.17 | 1.04 | 1.24                 |
| EASTERN<br>CANADA <sup>c</sup>      | 1.14                    | 1.47 | 1.22 | 1.59 | 1.19 | 1.38 | 1.36 | 1.37 | 1.60 | 1.39 | 1.66 | 1.80 | 1.18 | 1.87 | 1.55 | 1.74 | 1.76 | 1.53                 |

<sup>a</sup> Annual recovery unit productivity is an average for all pairs breeding in the recovery unit in the pertinent year.

<sup>b</sup> Unweighted average annual productivity from 1989 to 2023 for all states except New Hampshire (1997-2023).

<sup>c</sup> Includes productivity on the French islands of St. Pierre and Miquelon, reported by the Canadian Wildlife Service.




Figure 1. Abundance of Atlantic Coast piping plover breeding pairs by recovery unit, 1986-2023. Blue bars denote the annual pair estimate. Dashed pink lines indicate abundance objectives established in the 1996 revised recovery plan.